




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 4 : C12N 15/00, 7/00, 5/00		A1	(11) International Publication Number: WO 89/12684
			(43) International Publication Date: 28 December 1989 (28.12.89)
(21) International Application Number: PCT/GB89/00698		56 Bateman Street, Cambridge CB2 1LR (GB).	
(22) International Filing Date: 22 June 1989 (22.06.89)		(74) Agent: PERCY, Richard, Keith; Patent Department, National Research Development Corporation, 101 Newington Causeway, London SE1 6BU (GB).	
(30) Priority data:		(81) Designated States: AU, DK, JP, US.	
8815098.2 24 June 1988 (24.06.88) GB			
8904411.9 27 February 1989 (27.02.89) GB			
8907374.6 31 March 1989 (31.03.89) GB			
8910520.9 8 May 1989 (08.05.89) GB			
(71) Applicant (for all designated States except US): NATIONAL RESEARCH DEVELOPMENT CORPORATION [GB/GB]; 101 Newington Causeway, London SE1 6BU (GB).		Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.	
(72) Inventors; and (75) Inventors/Applicants (for US only): BINNS, Matthew, McKinley [GB/GB]; Appleby Cottage, Earith Road, Colne, Huntingdon, Cambridgeshire PE17 3NJ (GB). BOURSNEILL, Michael, Edward, Griffith [GB/GB]; 2 Lowry Close, St. Ives, Huntingdon, Cambridgeshire PE17 6EF (GB). CAMPBELL, Joan, Iyabo, Amiemenghena [NG/GB]; 35 Shakespeare Road, St. Ives, Huntingdon, Cambridgeshire PE17 4TT (GB). TOMLEY, Fiona, Margaret [GB/GB];			

(54) Title: FOWLPOX VIRUS NON-ESSENTIAL REGIONS

(57) Abstract

The invention relates to DNA from fowlpox virus (FPV) providing a non-essential region for the insertion of foreign genes therein and thence the construction of a vector for homologous recombination with a wild type FPV, whereby the resulting recombinant FPV can be used for vaccination of animals, especially chickens. In this invention, the non-essential region consists substantially of a length of DNA from the long unique sequence of the terminal inverted repeat (TIR) of FPV or from the region at FPV which corresponds approximately to the *HindIII* D fragment genes D8 and D9 in vaccinia virus.



FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FI	Finland	ML	Mali
AU	Australia	FR	France	MR	Mauritania
BB	Barbados	GA	Gabon	MW	Malawi
BE	Belgium	GB	United Kingdom	NL	Netherlands
BF	Burkina Fasso	HU	Hungary	NO	Norway
BG	Bulgaria	IT	Italy	RO	Romania
BJ	Benin	JP	Japan	SD	Sudan
BR	Brazil	KP	Democratic People's Republic of Korea	SE	Sweden
CF	Central African Republic	KR	Republic of Korea	SN	Senegal
CG	Congo	LJ	Liechtenstein	SU	Soviet Union
CH	Switzerland	LK	Sri Lanka	TD	Chad
CM	Cameroon	LU	Luxembourg	TG	Togo
DE	Germany, Federal Republic of	MC	Monaco	US	United States of America
DK	Denmark	MG	Madagascar		
ES	Spain				

- 1 -

FOWLPOX VIRUS NON-ESSENTIAL REGIONSBackground of the invention1. Field of the invention

This invention is in the field of recombinant DNA technology and relates to fowlpox virus as a vector for foreign DNA.

05 2. Description of the prior art

Several viruses with DNA genomes have been used to carry and express genes from other viruses or other species. Viruses used in such a way are known as 'vectors' and genes, other than their own, expressed in such a way are referred to as 'foreign genes'.

10 One of the primary requirements for a virus to be used as a vector in this manner is a suitable site for insertion of the foreign gene. If insertion of a gene into a site in the virus causes disruption of some function essential for growth, then such a site could not be used. Suitable sites are those at which
15 an insertion does not disrupt any functions or those whose functions are not essential for viral growth and therefore can be disrupted with impunity. Such sites are known as 'non-essential regions'. The phrase 'non-essential' in this context means non-essential for growth under at least some conditions in which
20 the virus can be grown in vitro and under at least some conditions in which it survives in vivo.

Examples of viruses which have been used as vectors by virtue of the fact that they contain non-essential regions are orthopoxviruses, adenoviruses and herpesviruses, although the
25 regions used may be different in each case. Vaccinia virus (VV), which has been used as a vector, is an orthopoxvirus, a member of the pox virus family. Fowlpoxvirus (FPV), the subject of this patent application, is also a pox virus, but is a member of a different genus, the avipoxviruses. VV can be grown in tissue
30 culture, and foreign genes can be inserted into the viral genome during this process. Several regions of VV have been found to be non-essential in vitro in more than one tissue-culture system. These include: large regions towards the left hand end, B. Moss et al., J. Virology 40, 387-395, (1981) who describe a mutant VV

- 2 -

having a deletion 6.4 megaDaltons from the left-hand end; D. Panicali *et al.*, J. Virology 37 1000-1010, 1981) who describe a mutant VV having a deletion starting 6.85 megaDaltons from the left-hand end; the thymidine kinase (TK) gene, D. Panicali and E. Paoletti, Proc. Natl. Acad. Sci. USA 79, 4927-4931 (1982); M. Mackett *et al.*, J. Gen. Virol. 45, 683-701, (1982); and the vaccinia growth factor (VGF) gene, R.M.L. Buller *et al.*, J. Virology 62, 866-874 (1988). Sites such as these might also be non-essential for growth *in vivo*. However in the case of both the TK and VGF gene, it has been found that although growth in tissue culture is unaffected or only slightly affected by insertion into the gene, growth and virulence *in vivo* are markedly affected, R.M.L. Buller *et al.*, Nature 317, 813-815 (1985) and *loc. cit.* (1988), showing that these genes are not completely non-essential for growth of the virus *in vivo*. This *in vivo* attenuation may however be useful if it reduces unwanted pathogenic effects of the virus and accordingly growth *in vivo* with accompanying attenuation is a valid growth condition for the purpose of this invention.

Some sites are essential in some tissue culture systems and non-essential in others. For example there is a gene in VV which is essential for replication in human cells but which is non-essential in chicken embryo fibroblast cells, S. Gillard *et al.*, Proc. Natl. Acad. Sci. USA 83, 5573-5577 (1986). Another gene in the related orthopoxvirus cowpox virus is essential for growth in chinese hamster ovary cells but not for growth in chicken embryo fibroblast cells, D. Spehner *et al.*, J. Virology 62, 1297-1304 (1988). These examples show that differences in the tissue culture systems can affect which regions are non-essential.

The VV genome is far from being completely mapped and relatively little has been published about the FPV genome. It is known that FPV has a TK gene which has about 60% homology with the VV TK gene at the amino acid level, D.B. Boyle *et al.*, Virology 156, 355-365 (1987). Like the VV TK gene, it serves as

- 4 -

end and that the deletion mutant virus is viable in tissue culture (cell type not clearly stated).

Some information has been released about the very terminal region of FPV, in a poster by J.I.A. Campbell displayed at the
05 International Poxvirus Workshop, Cold Spring Harbor, New York, 9-13 September 1987. (The poster gives more information than does the abstract published in "Modern Approaches to New Vaccines including Prevention of AIDS", CSH 1987, page 45). Like VV and cowpox virus, FPV has terminal sequences which are inverted
10 repeats of each other and are covalently cross-linked together. A part of the terminal inverted repeat (TIR) sequence was digested with BamHI and cloned. The BamHI fragment was described as having a length of 6.3 kb. The poster describes the general layout of the sequences, which from left to right comprise a
15 short unique region, a set of tandemly repeated sequences and then a long unique region containing three possible open reading frames. The summary notes that this FPV terminal fragment shares the general pattern of nucleotide sequence with the terminal fragments of VV and cowpoxvirus, but also notes some marked
20 differences. The sequence of part of the VV TIR, namely from approximately nucleotides 6800 to 9000 from the left-hand end, VV is reported by S. Venkatesan et al., J. Virology 44, 637-646 (1982).

Large differences between the DNA sequence of VV and FPV are
25 to be expected, since the FPV genome is estimated to be at least one third longer than that of VV. It seems likely, from the present knowledge as cited above, that many of the differences between the genomes of FPV and VV will be nearer to the termini than to the centre of the genome. Thus, information about the
30 terminal region of VV is of limited interest in relation to FPV.

It has been a problem to locate a well-defined non-essential region other than the TK gene, in fowlpox virus. Very recently (20th April 1989), in PCT Application Publication No. WO89/03429 (Health Research Inc.), FPV recombinants have been described, but
35 the non-essential regions mentioned therein are not all well

- 3 -

a non-essential region for homologous recombination under at least some conditions : see PCT Application WO 88/02022 (CSIRO). Using the E. coli xanthine-guanine phosphoribosyl transferase (Ecogpt) gene as a dominant selectable marker, in conjunction
05 with the VV "7.5" promoter, the influenza haemagglutinin (HA) gene was inserted into a TK region of FPV and the FPV grown in chicken embryo skin cells. Expression of the HA gene was demonstrated by the binding of HA antibodies and labelled protein A to plaques of recombinant FPV.

10 The genome of FPV is known to have other similarities to VV in some regions. For example the DNA polymerase genes are closely related, having 42% homology at the amino acid level, M.M. Binns et al., Nucleic Acids Research 15, 6563-6573 (1987). On the other hand, F.M. Tomley, in a talk at the 6th Workshop on
15 Poxvirus/Iridovirus, Cold Spring Harbor, New York, 24-28 September 1986, described an attempted correlation of an 11.2 kb fragment of FPV DNA, located near one end of the genome, with VV DNA. While she reported 25% amino acid homology between a gene predicting a 48 kd polypeptide in FPV and a gene in VV
20 coding for a 42 kd polypeptide, no other match of any significance was found. More detail of this work is given in F.M. Tomley et al., J. Gen. Virol. 69, 1025-1040 (1988).

F.M. Tomley et al., quoting M. Mackett and L.C. Archard, J. General Virology 45, 683-701 (1979), also state at page 1038 that
25 the terminal portions of orthopoxvirus genomes are less well conserved than the central region and also say that the terminal regions, of around 30 to 35 kb in length, are thought to encode genes which determine specific virus characteristics such as host range, virulence, tissue tropism and cytopathogenicity and are
30 thus less likely to be conserved.

The terminal region of VV is known to be of possible interest in relation to non-essential regions. Thus, M.E. Perkus et al., Virology 152, 285-297 (1986), have found that deletions in the VV genome occur in the presence of bromoxydeuridine, one of which
35 extended from about nucleotides 2700 to 24100 from the left-hand

- 5 -

defined. The recombinants were generated merely by cleaving the FPV gene into fragments and trying the fragments with test constructs to see whether the foreign gene under test was expressed.

05 The VV HindIII-D fragment is one of the best characterised regions of the virus. The sequence of this 16060bp fragment has been determined, Niles et al., Virology 153, 96-112 (1986), and thirteen genes (D1-D13) have been identified.

Several temperature-sensitive mutants have been mapped to
10 genes within HindIII-D and fine mapping has assigned these to D2, D3, D5, D6, D7, D11 and D13, Seto et al., Virology 160 110-119 (1987). These genes are therefore essential for virus replication. It is not known whether the remaining genes within the VV HindIII-D fragment are essential or non-essential for
15 virus growth. Recent experiments have suggested (though not proved) that D8, which encodes a transmembrane protein, might be non-essential for propagation of vaccinia virus in tissue culture, Niles and Seto, Journal of Virology 62 3772-3778 (1988). In these experiments, a frameshift mutation was
20 introduced into the carboxy-end of D8 which removed the carboxy-terminal 56 amino acids of D8. Virus containing this mutation had growth rates indistinguishable from those of wild type virus.

It was not predictable, however, whether the D8 gene would
25 occur in fowlpox virus and if so whether it would be non-essential. Moreover, there was the problem of how to locate the D8 gene in the FPV genome which is relatively unmapped and is much larger than that of VV. Consequently, examination of the vaccinia Virus HindIII-D fragment did not indicate how to find
30 further non-essential regions within FPV.

Summary of the invention

It has now been found that the three open-reading frames (ORFs) wholly within the above-mentioned BamHI fragment of the terminal inverted repeat (TIR) are non-essential regions which
35 can serve as sites for homologous recombination events whereby a

- 6 -

foreign gene can be introduced into FPV (refer to Figure 1). These ORFs are characterised in that they have lengths of approximately 670, 370 and 300 nucleotides and fall within the long unique sequence (LUS) of the TIR within the BamHI fragment.

05 The BamHI fragment has been found to have a length of about 6.3kb and the long unique sequence begins at approximately nucleotide 4100, numbering from the left-hand end. (As to nucleotide numbering, please refer to the cautionary note immediately preceding the DNA sequence shown hereinafter). The lengths of

10 the ORFs can also be expressed in terms of the approximate molecular weights of the polypeptides which they are predicted to encode, namely 26.0, 13.9 and 11.5 kilodaltons. This nomenclature will be used hereinafter for identification purposes. The 26.0 and 11.5 genes are coded for on the same

15 strand and in that order when reading the fragment left to right from the 5' end towards the BamHI site, the 11.5 gene having its 3' near the BamHI site (see Fig. 3). The 13.9 gene is encoded on the complementary strand, reading right to left, but within a region distinct from and lying in between that coding for the

20 26.0. and 11.5 genes on the first strand.

Extrapolation from this and other findings leads to the reasonable suppositions that the whole of the long unique sequence of the TIR is a non-essential region, including sequence lying to the immediate right-hand end of the BamHI site, i.e.

25 within the 11.2kb fragment sequenced by F.M. Tomley et al., loc. cit. (1988). The TIR has a length of about 10 kb, i.e. $10 - 6.3 = 3.7$ kb lies within the 11.2 kb fragment. Thus, the long unique sequence extends to the right from about nucleotide 4100 over the remaining lengths of TIR, of about 6 kb.

30 Accordingly, in an important aspect the present invention provides a recombination vector for use in producing a recombination fowlpox virus (FPV) by homologous DNA recombination, said recombination vector comprising a cloning vector containing a non-essential region (NER) sequence

35 comprising at least part of the long unique sequence (LUS) of the

- 7 -

terminal inverted repeat (TIR) of FPV, said NER being interrupted by a foreign gene or genes and by a sequence or sequences for regulating the expression of each foreign gene present.

Preferably the NER is within part of the long unique sequence (LUS) lying within the 6.3 kb BamHI fragment. In a more particular aspect, the NER consists of at least a homologously recombinable length of an open reading frame (ORF) having a length of approximately 670, 370 or 300 nucleotides and falling wholly within that part of the LUS within the BamHI fragment of the TIR. However, we have also found that some isolates of the same strain of FPV do not have a lengthy sequence of over 200 bp of DNA spanning the BamHI site, which is an important pointer to the non-essentiality of the sequence spanning that site. It is therefore inappropriate to use the terminology "BamHI fragment" in a formal definition applicable to FPV generally. Accordingly, these ORFs are better defined as lying within the first 2.2 kb of the LUS after the tandem repeats, that is to say within 2.2 kb of its external end. NERs within this 2.2 kb are preferred. The above preferences for location of NERs apply also to the flanking sequences contained in the recombination vector and this invention, i.e. all the homologously recombinable sequence which flanks the foreign gene(s) and their regulatory sequence(s).

We have also found certain other non-essential regions of fowlpox virus. Whereas those described above are within part of the terminal inverted repeat, these further regions are in a more central part of the FPV DNA molecule, namely within or adjacent to one end of a 6.3 kb fragment of an EcoRI library of FPV, the genes of which show significant degrees of homology with certain genes of the vaccinia virus (VV) HindIII-D fragment.

This second aspect of the present invention has its origins in a random cloning of parts of the FPV genome, sequencing of the random clones and a search in all six reading frames for possible open reading frames which might predict genes. The sequences thus generated were compared with vaccinia virus (VV) protein sequences. It so happened that degrees of homology were noted to

- 8 -

some of the VV proteins D1-D13, including approximate matches to VV D1, D5, D9, D10 and D13. It was therefore decided provisionally to adopt the "D" terminology for the FPV genes which showed some degree of homology with the VV genes. An M13 clone of FPV matching VV D9 was used as a probe to identify clones within an EcoRI library of FPV. One such clone, designated pMB379, containing a 6.3kb insert, was sequenced. This clone contained part of the D9 gene and all of D10, D11, D12 and D13. (The last four genes showed 40%, 62%, 53% and 55% identity of amino acids encoded to their VV homologues). In order to clone D7 and D8 genes, a DraI library of FPV DNA cloned in the plasmid pUC13 SmaI was constructed. Colonies were probed with a prime-cut probe derived from an M13 clone designated M379-27S (which lies within the FPV D9 gene) and one positive colony, containing a plasmid designated pMB389, was studied further. pMB389 was found to contain the "carboxy"-end of the D6 gene, entire D7, D9, D10 genes and the "carboxy"-end of the D11 gene. (D11 runs in the reverse direction to D10, so the carboxy end is that nearer to D10. The "carboxy" end is that which is the 3'- end of the coding strand, corresponding to the carboxy end of the predicted polypeptide). There was no identifiable homologue of the VV D8 gene, although another gene was present between D7 and D9. This has been termed "D8" as a matter of convenience. In a particularly unexpected finding, the copy of D9 present in both pMB379 and pMB389 possessed frameshift mutations when compared with its VV counterpart, thus making it unlikely that D9 is expressed in FPV.

It has now been found that the D8 and D9 genes of FPV are non-essential genes which can be used for the insertion of foreign DNA. Since the intergenic regions can also be considered non-essential, it follows that in principle the whole of the region following the carboxy-end of D7 and up to the carboxy end of the D9 gene is a non-essential region (NER). (The genes D7, D8, D9 and D10 all run in the same direction). However, as will be explained hereinafter, the NER does not run right up to the

- 9 -

carboxy-end of the D9 gene, but stops short thereof in order to avoid possible interference with the D10 gene, which overlaps with the D9, and the promoter of the D10 gene. This NER will be referred to hereinafter for brevity as the D8-D9 NER.

05 Accordingly, the present invention further provides a recombination vector as set forth above but wherein the NER is such a carboxy end-D7 to near carboxy end-D9 region NER (but excepting the D10 gene and D10 promoter region).

10 In a second important aspect the invention provides a recombinant vector, e.g. a plasmid, containing a NER sequence (TIR or D8-D9) as defined above. Such a vector is useful for preparing the above-defined recombination vector.

15 Thirdly, the invention includes a DNA molecule which consists substantially or essentially of a NER sequence as defined above with or without some homologously recombinable flanking sequence, say (arbitrarily) up to 1000 bp per end at either or both ends thereof in order to distinguish such a molecule from the genomic DNA of FPV.

20 Fourthly, the invention covers a host, appropriate to the vector, harbouring any of the vectors of the invention, i.e. the recombination vector or any intermediate vector which contains the novel NER sequence.

25 Fifthly, the invention embraces a recombinant FPV produced by homologous recombination of a parent FPV with the insert DNA of a recombination vector defined above.

30 Sixthly, the invention includes animal, especially chicken, cells infected with the above-identified recombinant FPV, a process of in vitro culture of said cells and, where patent law permits, a method of vaccinating a responsive animal, especially a chicken, with the recombinant FPV.

In a more particular aspect the recombination vector of the invention comprises in order:

- (1) a first homologously recombinable flanking sequence,
 - (2) a sequence within a first portion of the non-essential
- 35 region,

- 10 -

- (3) promoter DNA,
(4) a foreign gene transcribably downstream of the promoter (whereby when the fowlpox virus RNA polymerase binds to the promoter it will transcribe the foreign gene into mRNA),
05 (5) a sequence within a second portion of the same non-essential region, the first and second sequences being in the same relative orientation as are the first and second portions of the non-essential region within the viral genome, and
10 (6) a second homologously recombinable flanking sequence.

Brief description of the drawings

Fig. 1 is a schematic illustration of a homologous recombination process of the invention;

15 Fig. 2 is a map of the BamHI terminal fragment of FPV DNA showing the location therein of the beginning of the long unique sequence of the TIR;

Fig. 3 is a DNA map of the FPV TIR showing three non-essential region sequences useful in the present invention and other ORFs, together with a map of the VV TIR showing the
20 different ORFs found in VV;

Fig. 4 to 7 illustrate schematically various general methods of construction of recombination plasmids in accordance with the invention; and

25 Fig. 8 illustrates schematically the construction of specific recombination plasmids of the invention containing non-essential region sequence of FPV into which a foreign gene is inserted.

Figure 9 is a map of the genes D6 to D13 and an adjacent gene A2 showing the genes cloned in plasmids pMB379 and 389 and important restriction sites in these plasmids;

30 and

Figure 10 illustrates schematically the construction of specific recombination plasmids of the invention containing sequence within the D8, D9 and D10 genes (D8, D9 according to the invention; D10 comparative) into which a foreign gene is
35 inserted.

- 12 -

Below is shown the DNA sequence of the BamHI fragment, starting from the left-hand end and reading in the usual 5' to 3' direction, together with predicted polypeptides encoded. It should be borne in mind, however, that the sequencing of the

05 tandemly repeated region has presented difficulties and that the number of nucleotides within this region might be subject to some revision. This is not a matter of any significance to the present invention, which is not concerned with the tandemly repeated region. It follows that the long unique sequence does

10 not necessarily start at about nucleotide 4110 in absolute numbering and the nucleotide numbers are therefore to be treated as merely guidance to the reader to enable him to identify the ORFs referred to and to be construed accordingly in the description and claims.

- 11 -

Description of the preferred embodiments

Referring first to the terminal invected repeat (TIR) NERs and to Figure 2, the BamHI fragment of the TIR of FPV has a length of about 6.3 kb. It contains a short unique sequence of length 0.23 kb, followed by a region of approximate length 3.87 kb which contains tandemly repeated sequences and then a long unique sequence of length 2.18 kb.

The approximate location of the genes providing the TIR NER DNA for use in the invention within the BamHI fragment is shown in the map of Figure 3. The arrows show that the 13.9 gene is on a complementary strand. From this map and from sequence information it will be seen that these genes have no known near counterpart in VV. The BamHI site is believed to be part of a further non-essential region for the reasons explained above. A total of five ORFs have been found in the long unique sequence of the TIR. Those which appear to encode polypeptides by virtue of their distinctive FPV codon bias occur as follows :-

<u>Location (Nucleotides numbered from left-hand end)</u>	<u>No. nucleotides</u>	<u>No. amino acids encoded</u>	<u>M.w. of polypeptide</u>	<u>Unique site</u>
4125 - 4790	666	222	26.0 kd	<u>Bgl</u> II
5261 - 5629C	369	123	13.9 kd	<u>Nco</u> I
5883 - 6185	303	101	11.5 kd	<u>Spe</u> I

C = occurs on the strand complementary to that shown in the nucleotide sequence below.

Besides the three ORFs designated 26.0, 13.9 and 11.5, there are two shorter ORFs on the complementary strand. Since, however, neither of these ORFs exhibit the required FPV codon bias it is unlikely that they code for genes.

- 13 -

1 ACCCTCATCTTACGATGAGTATTTATATAGTAAAAAAATGTATAAACAGTACCTTCCAA
61 AACCCCTAATATACACATTCTTTTACCCCTTAATTTGTTAAGGTGTAAAAATACCCCTATT
121 AAAATATATATTATTGTTTTAATAAAAAAAACCATACGGTTTTACATAAAATAATACTAT
181 ATCTAATTTCTTCCGGAATAATTTTATAAAGCTACCCAACGTTAGCGAAAACTTTTT
241 TATCGAGAGCTCGAGTTATACAAAAAGTTTTTATCGAGATTTGAAAAGCTTTTTTATCG
301 AGAGCTCGAGTTATACAAAACTTTTTTATCGAGAGCTAGAGTTATAGAAAAAGTTTTTA
361 TCGAGATTTGAAAAGCTTTTTTATCGAGCGCTCGAGTTATAGAAAACTTTT.....

..... continuation of combinations of blocks of repeated
sequences.....

3721 .ATAGAAAAAGTTTTTATCGAGATTTGAAAAGCTTTTTTATCGAGAGCTCGAGTTATAG
3781 AAAAAGTTTTTATCGAGAGCTCGAGTTATAGAAAAAGTTTTTATCGAGATTTGAAAAG
3841 CTTTTTATCGAGAGCTCGAGTTATAGAAAACTTTTTTATCGAGAGCTCGAGTTATAGA
3901 AAAAGTTTTTATCGAGATTTGAAAAGCTTTTTTATCGAGAGCTCGAGTTATAGAAAA
3961 GTTTTTTATCGAGAGCTCGAGTTATAGAAAACTTTTTTATCGAGAGCTCGAGTTATAGA
4021 AAAAGTTTTTATCGAGAGCTCGAGTTATAGAAAACTTTTTTATCGAGATTTGAAAAG

↓(4101)

M I V E K I

4081 CTTTTTTTATCGAGAGCTCGAGAAGTTAAATCGAGACGCCGATATGATCGTAGAGAAAA
A A W L L Y P L C L L R C F L C N S V R
4141 TAGCGGCGTGGTTATTGTATCCGCTATGCCTTCTACGATGTTTCTCTGTAACGCGTAA
P A T C K C V H C L L Y P F E V C C E C
4201 GGCCCGCCACTTGCAAATGCGTCCACTGTCTTGTATCCCTTGAAGTATGTTGCGAAT
M S E T L D S L E H S C C Y C C V L P L
4261 GCATGAGCGAGACGTTAGACTCTCTGGAACACAGTTGTTGTTACTGTTGCGTGCTTCTC
L I I R E F W R R V I L P T L K A T C D
4321 TATTGATTATCAGAGAGTTCTGGAGACGCGTGATATTACCTACTCTAAAAGCGACTTGCG
C I R L P C V L T R R F C K R T I C P L
4381 ACTGTATTAGGCTACCGTGCGTTCTCACCAGAAGATTCTGTAAAAGAACCATCTGCCCGT
A K S W C R C F C C P C E V F L R C L L
4441 TAGCTAAATCTTGGTGTCGTTGTTTCTGTTGCCCTTGCGAGGTTTTCTTGAGGTGTCTCC
F P C M M L R R M H R G R L T G V R E P
4501 TCTTTCCTTGCATGATGTTGAGAAGAATGCATAGAGGCAGACTGACCGGAGTAAGAGAAC

- 14 -

G A F R D S R D P A R R G T W V N D W C
4561 CGGGAGCGTTTAGAGATTCAAGAGATCCCGCCAGACGGGGAACCTGGGTCAACGACTGGT

Bgl II
E D L C V W I W S P C C Y V K R C I R T
4621 CGGAAGATCTCTGCGTATGGATATGGTCTCCGTGTTGTTACGTCAAGAGATGTATTCGCA

M C D T F T K K I F Y W F I A P A G S P
4681 CGATGTGCGACACCTTCACTAAAAAATTTTCTACTGGTTCATCGCCCCGCAGGATCGC

R M P E E P S P L S R K V F S S *
4741 CGAGAATGCCCCGAGGAACCTTCTCCGCTATCGAGAAAGGTCTTTTCGTCTGAAGCATCG

4801 CGACGTGTAGATGAAAGAGATTGGACGCCAAAAGGTATACGCTCGTATTATTACGAATCG

4861 CGCTCGTCGTCTGCCTCGAAGATGATAAAGAGAACGCGAGTAAGGAAATACAAGATAAA

4921 AACAACTGTGATTCTACGTAATACGATTTCTAAAACGGCGTCTACCGCAATCTTTTTC

4981 ATCAGAGTAAGGGATAGGTTAAGATTCTTAAGGACCTTTATCAAACTATCGGAGAAAAA

5041 CTTTACGCTAAAGCCGCTCTCATGCTCTTTCACAGACTGGTGACCTATAGGCTTCCTATC

5101 GTTAGGGATATAGTCCGTTTTATTACGCGAGGAAAAATTTTGGAGGTTGCTTTTCTA

5161 TATCTCATGAAGAAAGTCGCTGTTTCGTCCTACTAAAACTAATGCTAAAAAAAATCTAT

5221 CGTTAATAAATTAAGTTATCGGGTTTTGAATATTATTAATCTATGCATACGAAACCGT
* D I C V F G Y

5281 ACGTCGCGTTACACGGGGAGAATCCGAACGTTTTCCCGTCCGTATAGGCACACTTACCCA
T A N C P S F G F T K G D T Y A C K G V

5341 CTTCTGGTTTCTTGTCATCCCTTATCCACAACTCTGCCCCATACCGAATTTGCTTATGA
E P K K D D R I W L S Q G M G F K S I F

Nco I
5401 ATTTTCATGTCTTTTTTATTGTCTATTAGACTACGTCTAGCTAACGTACCCATGGGAAACG

K M D K E N D I L S R R A L T G M P F T

5461 TTATCAAACACGTAGTCGACGCTTCGCTGTAAGTTTTTTTGTGATGCATTTTATCTAATC
I L C T T S A E S Y T K K H H M K D L R

5521 GGTTTACCCAGATACACTTATTGTTGTAAGAGGTGTACGGTCCTCTGCAATTTAGAGAGT
N V W I C K N N Y S T Y P G R C N L S N

5581 TAGCGCAATCCAACATTAACAACATCGCTATCAAGATTAATGATTTTCATTTTGATACACG
A C D L M L L M A I L I L S K M

5641 GATCAATTTTAATAGTCTATAGAGATATACCCACCAATAGTTACGCGATTAATTTTCAA

- 15 -

5701 TTATACATAATAAAATCTAGAGGAGTTGCCATGTGATTCTCTGTCGACGACGTTACCTC

5761 GTTAGTTTTATTAATTCTTCTTTAAGTTCTGACGACGGTTATATACCAAACAGCTTTTA

5821 CTTTTCGTAAGACGATAATTACGTTATAAAAGATTAAGAAATTTTATTTTACTAAAGTA

5881 M N N D T I F T L F Y C K N K K Y V R G
TCATGAATAATGACACGATATTTACTTTGTTTTATTGTAAAAATAAAAAATATGTTTCGTG5941 E G G R R R G K T G I L L F H P I N H R
GGGAGGGGGAAGGAGGAGAGGGAAGACGGGTATCCTTTTATTCCATCCAATAAACACCSpe I
6001 V I G T S A H Q C Y K T R R I G F K L Y
GTGTTATAGGAAGTGTGCACATCAGTGTTATAAACACGAAGAATCGGTTTCAAACCTCT
-----6061 A V A P R H V S T I R C G R S H S A H R
ACGCAGTTGCCCCTCGTCACGTATCCACTATCCGCTGTGGGAGATCCCACTCGGCTCACC6121 V D K F S F S F Q K V D F H C I A G S G
GAGTAGATAAGTTCTCCTTCTCCTTCCAAAAGTAGATTTTCACTGTATCGCCGGCAGCGA *
6181 GCGCATAGGTATCTGGCCCTACAGAATTCATATTCCTGATAGCTCGCGCGACGAAGCCT

6241 CCGTCGCCCCTGTGATCATTTTCATAATTCAAGGGGATCC (6280)

- 16 -

It will be appreciated that the sequence shown above is not definitive of FPV generically, since there will inevitably be small variations between different types or strains of FPV, some of which might take the form of deletions or insertions, and any
05 variant NER compatible with the desired homologous recombination is usable herein. In particular, DNA having a NER sequence based on ORFs having a sequence showing at least, say, 90% homology with the sequence set forth above is a preferred feature of the invention.

10 Beyond the BamHI site (GGATCC) the nucleotide sequence continues as per F.M. Tomley et al. loc. cit. (1988) at page 1029, nucleotide 6275 shown above being the first of the BamHI recognition sequence and No. 1 in Tomley et al.'s numbering. While there are no ORFs spanning the BamHI site, the sequence
15 extending at least as far as the first 160 bp into the (11.2 kb fragment of Tomley et al. is demonstrable non-essential, while further sequence thereafter is likely to be.

The NERs of the invention can be used to insert foreign genes into FPV. It is desirable that the gene be stably incorporated
20 so that the resultant recombinant FPV can be repeatedly cultured in tissue culture on a large scale. Evidence to date is that the 13.9 gene is less stable than the other two.

The foreign gene can be inserted at any location within the NER sequence of the recombination plasmid. Preferably it is not
25 so excessively near one end or the other as not to provide a site long enough for homologous recombination with the corresponding NER of the wild type FPV. However, it is possible to allow homologous recombination to occur outside the NER by providing, in the recombination vector, flanking sequence lying beyond
30 either or both ends of the NER. It is probably not usually beneficial to use flanking sequences which are heavily tandemly repeated, since this will increase the chances of unwanted recombination events. Thus, it is suggested that flanking sequences running extensively beyond the external end of the LUS
35 into the tandem repeat region are best avoided.

- 17 -

Referring now to the D8-D9 NER and to Fig. 9 of the drawings, the plasmids pMB379 and pMB389 have FPV DNA inserts which overlap in the D9 and D10 gene region. By sequencing these inserts a long nucleotide sequence has been obtained which runs from the 3'-end of D6 through to D13 and beyond that to the adjacent A2 gene.

In the sequence shown below, those parts which are not relevant to the present invention have been truncated. The restriction sites DraI (TTTAAA), EcoRI (GAATTC), BclI (TGATCA), AsuII (TTCGAA) and BglII (AGATCT) used in the cloning and construction of plasmids for testing out non-essential regions are underlined. The amino acid sequences of predicted genes are given in the single letter code. The D9 gene, being considered to be unexpressed in FPV, is not given an amino acid sequence, but is considered to run from nucleotides 1298 to 1984. Start and stop codons have been underlined. The sequence shown encompasses the carboxy end of the D7 gene and the carboxy end of the D11 gene. (Since the D11 gene has an open-reading frame running in the reverse direction to the others, its near-D10 end is that encoding the carboxy terminus of the putative D11 protein). The sequence shows the DraI site in the D11 gene (see also Figure 9).

It will be seen that the D10 gene starts within the end of the D9 gene, being within a different reading frame. The D10 promoter is therefore also within the D9 gene. The non-essential region is therefore defined as extending up to the beginning of the D10 promoter region, which is estimated at approximately 35 bases before the ATG start codon for D10, i.e. at around nucleotide 1936.

- 18 -

D7 (carboxy-terminus)

A Q S G M E S S F V F L G N I I E K *
1 AGCCCAATCTGGAATGGAATCATCTTTTGTCTTCTAGGTAACATAATTGAAAAATAATA
61 CTTTAATAAAGATTAATTTACATAATACCTACATCATGTAACGTATAGCTTATTAAACAT
121 GGATAGCATTTAGTAATATTTGTAAAATCAAAAATAACTATAAATATTTTAATAGTATAT
181 TGAATTCTTTATCATGTAGAAAGATATTAGTTATACACACAATGATATGTAAAAATAGTA
241 TTCGACATAAAGAATTGACGTTGATAATATTTCTCTGCTTTATATATCTATCAAAAACCT
301 TTGTAGAAGTAAAAGCACCTCCCTATGTACTCGTACCGGAAGTAGTGATATAAATCTTAC
361 TTGTTTTATTAAAGATGATCAGGGTGCTGGAGAAGATAAAGTAATTGTAGCGTGGCAACA
421 GGGTGACGGAGAAGTTACTCAAGGAATAAAAACAACTTGGGATACGACACGACAAGTGGG
D8 M L H I S K V S K E A E G S Y M C V
481 CCAAACCATGCTACATATATCCAAGGTATCTAAAGAAGCAGAAGGTCATATATGTGTGT
V W I N G D A D Y K K M N L G V F T V S
541 TGTATGGATAAACGGCGACGCTGATTATAAAAAAATGAATCTAGGAGTGTTTACTGTTTC
K K T Y M H N E V S I T P R R S R V D M
601 AAAAAAACCTACATGCATAACGAAGTAAGTATAACCCCTCGAAGATCGAGAGTAGATAT
S D G R P L K I E C A F S T R R I Y G R
661 GTCGGATGGTCGGCCGCTAAAAATTGAATGTGCTTTTAGTACCCGACGAATATATGGTAG
S Q V N I K W W K I D G I T R K W E Q Q
721 AAGCCAAGTCAACATAAAATGGTGAAGATTGACGGTATTACACGAAAATGGGAACAACA
T S G V N L L L H T Y G G I G S L S I P
781 GACTTCGGGAGTTAACTTACTGTACATACTTATGGAGGGATAGGATCGCTAAGTATCCC
N P T T G E S T G K Y M C V V T C G D I
841 AAATCCTACAACAGGAGAATCAACAGGTAAGTATATGTGCGTAGTTACCTGTGGAGATAT
G N V G F R L V K S L S P L S D T E S D
901 TGGAAATGTTGGATTTAGACTGGTAAATCATTATCTCCTTTATCAGATACAGAATCTGA
H S Y T S E E G S H F M E R C K V K K S
961 ICATAGTTATACTTCAGAGGAAGGATCACATTTTATGGAAAGATGCAAAGTCAAAAAAG
P Y G G W I V E *
1021 TCCGTACGGTGGATGGATAGTAGAATAGATCATCTCAGGGACGCGGTATATATTGCTATA
1081 AAGTAAAATAATTTCTCAATACATTTTCTACTTTAACGATATAATCGTTAACTTATTAG
1141 TTTAATTATATCATATATTCCTTTATATCTTAATTAATAAAAAATATATTCATAAATGGGAT
1201 AACTAGTATTTTTTTATATAGTTTACGATATGCTATTTCACTATATATAGTTTATCAAA

- 19 -

1261 CATAAGTAAAAATAAACAACTATAAAAACTAAGATCATGTTCGATATATCCAGAGAACA
1321 ACAAATATATTAGAAAAAAATAAAGATTGTGTTATTACTTTCGAAACAAATAAAGAATA
1381 ACTATAGAAAACACTAACATAAAAGATATACTTAGCGATAGACGAATTCACATATCGCGT
1441 TATGTATTACATCAGATAATATACCTATAATCGGTATAAGAAGAACGTCTTTTATGTATC
1501 AATCGGTTATATCAAAAAGAAGAAGCTTTTCAGAAATATTAGCCGTCGATATAAACCATC
1561 TAAAGTATATGTATAATAACGAAATTAAAGAAATATCATTAGATCAATAGTACCATTAC
1621 ATATAGCGGATTCAACAACCTTTGAAGAGTTAGTATTATTAGGAGGAAGAGTTAAAAATAA
1681 AGAATCAATATATCAATGCCTGAGTAGAGAACTATCGGAAGAAAGCGATGGAATACTTAC
1741 TATAAAAACATTTGGTAATAAAATATTAACCTTACCATAGAAGATAAGATACTTCGTAG
1801 AACATTTTATGGGTATTGTATAGTGTGTTTTATAGACCAACTCTATTCAGAAATCATTA
1861 ACCTTTATATAACATAGAAATCAAAGAATTAGGATCATTATTTGATCGATCAAGTAATGA
D10 M G E
1921 AAAATACGAATACTTGCATTTTATTTATAATACCTTATTAACATATAAATATGGGGGAGT
Y Y K N K L L L R P S V Y S D N I Q K I
1981 ATTATAAAAATAAATTGCTCCTTAGGCCGTCTGTATATTCTGATAATATACAAAAAATCA
K L V A Y E Y G K L H A K Y P L S V I G
2041 AACTGGTAGCATATGAATATGGCAAACACATGCTAAATATCCACTATCCGTAATAGGTA
I M K T I D D K F V L C H R Y N S F L F
2101 TAATGAAAACAATAGATGACAAGTTTGTGTTTTATGCCATAGATACAATAGTTTCTTGTTTT
S E I A F T K D K R R K I R L F K K Y S
2161 CAGAAATAGCATTTACAAAAGATAAACGACGGAATAAAGACTCTTTAAGAAATATTCAA
K Y M S N I E R D I L S Y K L S L P N N
2221 AGTATATGAGTAACATCGAACGTGATATATTAAGTTATAAGCTATCGCTTCCTAATAACT
Y N T N H I D I I F P G G K I K D L E S
2281 ATAATACAAACCATATAGATATAATCTTCCCAGGTGGTAAAATAAAGACTTGGAAAGTA
I T N C L V R E I K E E L N I D S S Y L
2341 TAACTAATTGTCTAGTAAGAGAAATAAAGAAGAATTAAATATCGATTCTCTTATCTCG
A I C K N C F V Y G S I Y D R L I D K D
2401 CTATCTGTAAGAACTGTTTTGTATACGGTCTATATACGACAGATTGATAGATAAAGATT
F E V I A L Y V E T D L T S R Q I L N R
2461 TTGAAGTTATAGCCCTTTACGTAGAAACAGATCTTACGAGTAGACAAATATTGAATAGAT

- 20 -

F I P N R E I K G I S F I D A R D I N K
2521 TCATCCCTAATAGAGAAATTAAAGGAATATCATTATAGACGCAAGAGATATTAACAAAG

D Y L Y T N V I K Y I I N A V R T S A S
2581 ATTATTTGTATACTAATGTAATTAAATATATAATAACGCTGTAAGAACATCCGCTAGCA

N S *
2641 ATAGTTAACATCTATCTAACTGTTAATATATAAATTAATT....
D11 * R V T L I Y I L K
(carboxy-terminus)

2881TAATTTTAAACCC.....
L K K F G

- 21 -

There are some preliminary indications that the insertion of a foreign gene in the D10 gene is unstable. Accordingly the D10 gene is not a NER for the purposes of the invention.

Referring now to all the NERs of this invention, the
05 recombination vector can contain a reasonable length of homologously recombinable flanking sequence to each side of the NER, preferably at least 100 nucleotides, more preferably at least 250 nucleotides and still more preferably at least 500
10 nucleotides. On the other hand, in the interests of not making the insert for the recombination vector too large, it is often desirable to limit such flanking sequence outside the NER to, say, not more than 1000 bp.

Incidentally, precise homology of the flanking sequence and the NER is not necessarily required. Remarks to that effect in
15 WO 89/03429 apply here also.

In preparing recombinant FPV of the invention, the first requirement is for a promoter. The well-known vaccinia virus P7.5 promoter the H6 promoter (see the cited PCT Application Publication No. WO 89/03429) or the P11 promoter, can be used, as
20 these are accepted by the FPV as compatible. Preferably, however, a FPV promoter is used. Certain promoters have been described in the prior PCT patent application Publication No. WO 89/03879 (National Research Development Corporation), the contents of which are herein incorporated by reference. The "4b"
25 and "13.2K" promoters therein are especially preferred.

The invention has particular relevance to poultry, especially chickens. For this purpose, any foreign gene relevant to improving the condition of poultry could be inserted into the fowlpox virus. Preferably the gene will be one appropriate to an
30 in vivo sub-unit vaccine, for example one or more genes selected from Infectious Bronchitis Virus (IBV), Infectious Bursal Disease virus, Newcastle Disease Virus (NDV), Marek's disease virus, infectious laryngotracheitis virus and genes encoding antigenic proteins of Eimeria species. Particular genes of interest are
35 the spike genes of IBV and the HN and F genes of NDV as described

- 22 -

in PCT Patent Application Publication No. WO 86/05806 and European Patent Application Publication No. 227414A (both National Research Development Corporation). In order for the foreign gene to be correctly translated in vivo it is necessary
05 for the foreign gene to have its own ATG start codon inserted in the region just following the promoter.

More than one foreign gene can be expressed in a single poxvirus. It is possible to arrange for mRNA for two foreign genes to be transcribed in different directions when inserted in
10 a single non-essential region. Such a "back to back" construction is shown in Example 7. Also, individual foreign genes can be inserted in individual NERs described herein. While fusion protein genes can, of course, be made, it is ordinarily preferable that each gene be under control of a separate
15 promoter.

The promoter and foreign gene then have to be inserted into the non-essential region (NER) of the FPV genome. The procedure of homologous recombination, illustrated by Figure 1 of the drawings, provides a way of doing so. A fragment of genomic DNA
20 containing the NER is sub-cloned in a cloning vector. A construct is then made, in the cloning vector, comprising part of the NER followed by the promoter followed by the foreign gene, followed by a further part of the NER in the same orientation as the first part. This construct, in an appropriate vector, forms
25 the recombination vector which is used to transfect the FPV, e.g. by the calcium phosphate method, whereby recombination occurs between the NER sequences in the recombination vector and the NER sequences in the FPV. The FPV then automatically re-packages this altered genome and the thus altered FPV (recombinant FPV) is
30 part of this invention.

The above-described recombination vector can be prepared in many different ways, the requisite elements being introduced in any order. Referring to Figure 4, a non-essential region defined above for use in the invention possessing (say) two restriction
35 sites A, B is inserted in an appropriate vector, which by way of

- 23 -

illustration only will be described as a plasmid. In another plasmid having the same (or ligatably compatible) restriction sites A, B, a construct is made of promoter sequence followed by the foreign gene sequence. It is of course essential that this
05 construct is made so that the mRNA transcription will begin at or before the start codon of the foreign gene. Since it is time-consuming to determine precisely where the mRNA transcription start is effected by any particular promoter, it is convenient simply to insert, say, 100 or more preferably 150 base
10 pairs of promoter DNA immediately preceding the FPV gene which it normally promotes, to ensure good working of the promoter.

The restriction sites A, B are located in the plasmid DNA flanking the FPV promoter DNA and the foreign gene. Of course, A can be within the promoter DNA if it falls within a
15 non-functional portion thereof. While A and B can be two different restriction sites, they can, of course, be one and the same. They can be sticky- or blunt-ended sites and can be prepared artificially, e.g. by filling in with additional nucleotides or chewing back, in ways well known in the
20 recombinant DNA field. Conveniently A and B are or are converted into a single blunt-ended site (C) and then allowed to ligate into a single blunt-ended site (C) within the NER. Care will have to be taken, of course, to select sites which are unique in the plasmid DNA to prevent ligation of other sequences of DNA
25 from occurring. In the Example herein, unique sites for SpeI, NcoI and BglII in the NERs have been used and have been filled in to provide a single blunt-ended site for insertion of the promoter-gene construct.

DNA from the two plasmids are ligated together in vitro and
30 then transformed into the host, to produce the final recombination plasmid of the invention.

Figure 5 illustrates another method of preparing recombinant vectors. In this method one first prepares a construct comprising a first part of the NER, followed by the promoter,
35 followed by a short sequence of nucleotides containing at least

- 24 -

one cloning site for introduction of a foreign gene, followed by a second part of the NER which will almost inevitably be in the same orientation as the first part. Of course, virtually any length of DNA would provide a cloning site suitable in some way or other for introducing a foreign gene. Preferably these constructs contain a multiple cloning site, that is to say a length of DNA containing the sites of a variety of different restriction enzymes, for example at least ten. Such a construct then has versatility, since it will then be much easier to restrict DNA flanking almost any foreign gene at sites close to each end thereof and insert the foreign gene into the multiple cloning site illustrated in Figure 5. Only two sites X, Y have been shown, for simplicity and these can be filled in or chewed back, as desired, to give identical blunt-ended sites (Z, replacing X, Y). In the final recombination plasmids the promoter DNA will be separated from the foreign gene by a portion of the multiple cloning site, but this will not adversely affect the transcription of the mRNA in the final virus.

In either method of construction the NER is split by the promoter and foreign gene. It is, of course, not essential that it be split in a central region. Nor is it essential that the second part thereof constitute the entire balance or remainder of the NER. As explained above, so long as there exists within or at each end of the NER contains a long enough stretch of DNA for homologous recombination, it does not matter that a part of the NER might be excised somewhere in between or that additional (irrelevant) DNA be inserted in preparing the recombination plasmid, or even that the NER is only a few bp in length (but it should not normally be less than about 20 bp long or there would be practical difficulties in inserting the foreign DNA).

Figure 6 illustrates another method of preparing a recombination plasmid. The NER plasmid is restricted as before (a blunt-ended site or sticky ended site filled in to become blunt-ended, C, is illustrated arbitrarily for this Figure). The other plasmid contains a marker gene with its own promoter and a foreign gene with its promoter in opposed orientations.

- 25 -

Different promoters should be used to avoid the possibility of an incorrect recombination. When these plasmids are ligated together the marker is linked to the foreign gene. The presence of the marker in the recombinant FPV is therefore likely to indicate a successful incorporation of the foreign gene into FPV. Conveniently the marker is readily detectable by a colour, fluorescent or chemiluminescent reaction. The beta-galactosidase or lacZ gene can be inserted into the NER and recombinants detected by the blue plaques generated when the 5-bromo-4-chloro-3-indolyl-D- galactopyranoside (X-gal) substrate is present in the growth medium. This technique, using the lacZ gene, has been described in relation to VV by S. Chakrabarti et al., Molecular and Cellular Biology 5, 3403-3409 (1985).

Preferably the marker gene is also selectable (or it is replaced by a selectable gene or two separate genes, one marker and one selectable are included). An example of a selectable gene is the Ecogpt gene described in PCT Application WO 88/02022, already cited, and also by F.G. Falkner and B. Moss, J. Virology 62, 1849-1854 (1988).

It is also useful when preparing a recombination plasmid to know that the foreign gene is likely to have been successfully inserted into the NER. One way of checking is to adopt the procedure shown in Figure 7. Again, blunt-ended sites (Z and W) are shown. In a first plasmid, containing the NER, a marker gene is inserted within the NER and optionally a multiple cloning site is inserted within the marker gene. The foreign gene with its promoter, contained in a separate plasmid, is then inserted into the marker gene by combining the two plasmids as shown. The foreign gene is inserted at any site within the marker gene which will interrupt or change its marker function, whereby the insertion can be recognised. For example, the E. coli. lacZ gene can be used as the marker gene. Only the alpha-fragment need be inserted since the remaining elements of the gene required for processing are provided by the bacterial host. It is provided with the E. coli. promoter for lacZ, the lacZ start codon and,

- 26 -

preferably, a multiple cloning site within a few codons of the start, followed by sequence encoding the alpha fragment. When the alpha-fragment is interrupted by a foreign gene, conveniently insertable in the said multiple cloning site, the blue colonies given on an X-gal substrate are not ordinarily obtained. White colonies indicate a likely recombination plasmid.

The recombination vector could, on occasion, be further provided with a termination signal site to terminate transcription of mRNA, which might be useful for stabilising the mRNA when an early-acting promoter is used. This would be inserted downstream of the foreign gene. However, in many cases the ordinary termination signals of the infecting poxvirus or the signals for starting transcription of another, downstream, gene of the FPV will be adequate.

References herein to vectors other than FPV (or VV) mean any convenient prokaryotic or eukaryotic cloning vector appropriate for bulk production of the construct within a suitable host. Prokaryotic vectors will ordinarily be plasmids or phages. Suitable prokaryotic hosts include bacteria. Eukaryotic vectors such as those of fungi, yeasts and animal cells, e.g. SV40, can be employed if thought more convenient.

Although the recombination vector used will ordinarily be of double-stranded DNA, it is possible to use single-stranded DNA for the homologous recombination.

The recombination vector of the invention containing the NER, promoter and foreign gene then has to be "swapped over" for the "parent" FPV DNA. For this purpose, appropriate animal, preferably but not necessarily poultry, cells are infected with the parent FPV, and the recombination vector is then introduced into the cells. This process is carried out in vitro. It is best not to use wild type parent FPV for obvious reasons. FPV can readily be attenuated (mutated to make it less virulent), by any conventional method of attenuation.

Many different methods are available for detecting or selecting the recombinant viruses, and have been described for VV

- 27 -

in a review article of M. Mackett and G.L. Smith, J. General Virology 67, 2067-2082 (1986). Such methods are applicable in the present invention. One method is to enlarge the recombination plasmid to include along with the desired foreign gene an additional marker gene as described above (Figure 6).
05 Alternatively, the insertion of the foreign gene can be detected by a hybridisation assay in which a labelled nucleotide sequence complementary to the foreign gene sequence is employed. Such assays are well known.

10 The FPV recombinants possessing the foreign gene are then grown in appropriate cells. The NER of the invention are non-essential in chicken embryo fibroblast cells. It would be necessary to carry out routine experiments to determine whether they are also usable for growth of FPV in other kinds of cell,
15 e.g. chicken embryo skin cells, chicken fibroblasts, chicken embryo epithelial cells derived by conventional tissue culture methods, principally trypsinisation of tissues or the chorioallantoic membrane (CAM) of embryonated chicken or turkey eggs.

20 For administration to birds as a vaccine, the recombinant virus can be given to birds by any suitable method such as by aerosol, drinking water, oral, intramuscular injection or inoculation into the wing web. Ingredients such as skimmed milk or glycerol can be used to stabilise the virus. It is preferred
25 to vaccinate chicks 1 day old by aerosol administration. A dose of from 10^2 to 10^8 pfu, preferably 10^4 to 10^6 pfu, of the recombinant FPV per bird is recommended in general.

While the invention is intended primarily for the treatment of chickens, it is potentially of interest in relation to the
30 vaccination of other animals which might safely be inoculated with FPV. It is even possible that it might be considered safe to infect humans with FPV after appropriate trials have taken place.

The following Examples illustrate the invention.

- 28 -

EXAMPLE 1

This Example relates to terminal inverted repeat NERs.

Virus strain

05 The HP438 strain of fowlpox virus was obtained from Professors A. Mayr and H. Mahnel, Ludwig-Maximilians University, Munich. The HP438 strain has been obtained from the pathogenic HP1 strain by 438 passages in chick embryo fibroblasts (CEFs) in tissue culture, A. Mayr et al., Zentralblatt für Veterinärmedizin B13: 1-13 (1966). The HP 444 strain used to obtain DNA for cloning was derived by 6 further passages in CEF cells.

10 Tissue culture medium

CEF cells were grown in 199 (Wellcome) medium, supplemented with Penicillin (200U/ml), Streptomycin (200µg/ml), Fungizone (2µg/ml) and 10% newborn calf serum.

Cell Culture and Viral DNA purification

15 FPV strain HP444 was grown in primary chick embryo fibroblast (CEF) cultures and FPV DNA prepared as described by M.M. Binns et al., Nucleic Acids Research 15, 6563-6573 (1987).

Identification of the terminal fragment

20 The genome of poxviruses is double-stranded DNA. As usual the two strands are held together by hydrogen-bonding. However, at the ends of the genome of poxviruses the two strands are covalently linked. Topologically this means that the DNA is a continuous strand. When the genome is cut into fragments with a restriction enzyme, internal fragments are topologically
25 different from terminal fragments and behave differently. On heating (to separate the two strands) the internal fragments completely separate and they reanneal slowly (hardly at all if one cools quickly). The terminal fragments, being covalently bonded, cannot separate very far spatially. On cooling they
30 reanneal quickly by starting at the middle and "zipping up". This is called 'snap-back'. After heating and rapid cooling therefore, only the terminal fragments can be seen by gel electrophoresis. This method of identification of the terminal fragment, called "the snap back method," see L.C. Archard & M.
35 Mackett, J. General Virol, 45, 61-63 (1979), was used here.

- 29 -

A BamHI restriction digest (10µg) DNA was extracted once with phenol, the aqueous phase was extracted with diethyl ether to remove phenol and the DNA in the aqueous phase was precipitated with ethanol. The DNA pellet was dried and resuspended in 20µl of TE (10mM Tris HCl pH 7.5, 1mM, EDTA). The DNA solution (10µl) was then denatured by the addition of 40µl of 95% v/v deionized formamide in 50mM Tris HCl pH 7.5 and incubated at 60°C for 10 minutes. The sample tube was then rapidly cooled on wet ice containing NaCl for 10 minutes before analysing the DNA by agarose gel electrophoresis. The presumed terminal fragment renatured rapidly only in the native FPV DNA and was detected as an approximately 6.3kb band. There is only one BamHI - derived snap back fragment, as the BamHI site falls well within the TIRs.

Cloning of the end fragment of FPV DNA

The terminal crosslinks of the BamHI fragment were digested with S1 nuclease; using 100µg of FPV DNA and 20 units S1 nuclease (Boehringer Mannheim) in 250µl of 30mM sodium acetate, pH 4.5; 0.3M NaCl; 1mM ZnSO₄; 5% (v/v) glycerol at 37°C for 20 minutes. To end-repair the DNA, it was then ethanol-precipitated and re-suspended in 100 µl of ligation buffer (50mM Tris. HCl pH 7.5, 10mM magnesium chloride, 10mM DTT) containing 5 units of E. coli polymerase I Klenow fragment and 1 unit of T₄ DNA polymerase and 0.025 mM each of dATP, dCTP, dGTP, and dTTP. This reaction was left for 1 hour at room temperature. It was stopped by heating for 10 minutes at 70°C and 20µl thereof was digested with 10 units of BamHI (Boehringer Mannheim) in accordance with the manufacturer's recommendations. 5µl of this DNA was ligated to 20 ng of BamHI/Sma I - cut plasmid pBGS19, B.G. Spratt et al., Gene 41, 337-342 (1986), to make a total volume of 20µl ligation buffer containing 2 units of T₄ DNA ligase and 1mM ATP, and incubated at 4°C overnight and transformed into E. coli. Screening for recombinants which contained the end fragment was carried out with a radiolabelled FPV BamHI terminal fragment isolated from a gel, using standard procedures as described by T. Maniatis et al. "Molecular Cloning: A Laboratory Manual",

- 30 -

New York, Cold Spring Harbor Laboratory (1982) and Holmes and Quigley, Analytical Biochemistry 114, 193-197. A recombinant plasmid pB3 was isolated.

Cloning of NERs

- 05 The three genes under test for non-essentiality were retained together in their natural sequence in a shortened modification of plasmid pB3, see Figure 8. pB3 was shortened by removing the short unique sequence and the repeated sequences (see Figure 8) from the BamHI fragment. This was done by a digestion with SstI
- 10 There is a SstI site in pBGS19 just before the BamHI site into which the FPV BamHI fragment was inserted. The SstI digestion was carried out at 37°C in a total of 20µl of the recommended buffer, using 10 units of SstI. There are many SstI sites in the repeated sequence, the last of which causes cutting at nucleotide
- 15 4005. Thus, after cutting with SstI, the plasmid pB3 (9.8kb) was re-ligated in a 100µl volume, to favour intramolecular ligation using the previously described ligation buffer, at 15°C overnight, to produce plasmid pB3ME (5.8kb) which lacks 4.0kb of the BamHI fragment DNA. To select the 5.8kb clones, rather than
- 20 those which have lost shorter lengths of SstI- cut DNA, selected colonies were grown up and then subjected to gel electrophoresis. To check that all the repeats were removed, the plasmid was digested with BglII and the fragments analysed by gel electrophoresis.
- 25 Preparation of a plasmid containing the VV 7.5K promoter and the lacZ gene

- The VV 7.5K promoter is available in a plasmid pGS20 from Dr. Geoff Smith, Division of Virology, Dept. of Pathology, University of Cambridge, Addenbrooke's Site, Hills Road, Cambridge,
- 30 England. Cambridge and is described in M. Mackett, G.L. Smith and B. Moss J. Virology 49, 857-864 (1984) and can be combined with E. coli. lacZ sequences as shown in the DNA sequence below. The resultant plasmid is designated pPG1. Other methods which could be used to combine the VV 7.5K promoter and lacZ sequence
- 35 have been described in papers relating to VV.

- 31 -

```

<----- p7.5 sequences from pGS20 -----
1  GGATCCCCAATTCCAGCTTGGCTGCAGGTCGACATATACTATATAGTAATACCAATACTC
-----
61  AAGACTACGAAACTGATACAATCTCTTATCATGTGGGTAATGTTCTCGATGTCGAATAGC
-----
121  CATATGCCGGTAGTTGCGATATACATAAACTGATCACTAATTCCAAACCCACCCGCTTTT
-----
181  TATAGTAAGTTTTTCACCCATAAATAATAAATACAATAATTAATTTCTCGTAAAAGTAGA
-----
-----p7.5 sequences from pGS20 ----->
241  AAATATATTCTAATTTATTGCACGGTAAGGAAGTAGAATCATAAAGAACGTGACGGATCG

<-SV40 sequences + linking DNA sequences -----><-- lacZ sequences
301  AAGCTTTGCAAAGATGGATAAAGTTTTCCGGAATTCCAAGCTTGGCCTGGCCGTCGTTTT
      MetAspLysValPheArgAsnSerLysLeuGlyLeuAlaValValLeu
-----
361  ACAACGTCGTGACTGGGAAAACCCTGGCGTTACCCAACCTTAATCGCCTTGACGACATCC
      GlnArgArgAspTrpGluAsnProGlyValThrGlnLeuAsnArgLeuAlaAlaHisPro
-----
.....continuation of lacZ sequences .....
-----
3301  GGGGATTGGTGGCGACGACTCCTGGAGCCCGTCAGTATCGGCGGAATTCCAGCTGAGCGC
      GlyIleGlyGlyAspAspSerTrpSerProSerValSerAlaGluPheGlnLeuSerAla
-----
----- lacZ sequences -----><--linking
3361  CGGTCGCTACCATTACCAGTTGGTCTGGTGTCAAAAATAATAAACCAGGGCAGGGGGGA
      GlyArgTyrHisTyrGlnLeuValTrpCysGlnLys***
-----
sequences--><----- SV40 sequences -----
3421  TCTGAACCTGTTTATTGCAGCTTATAATGGTTACAAATAAAGCAATAGCATCACAAATTT
-----
3481  CACAAATAAAGCATTTTTTTCACTGCATTCTAGTTGTGGTTTGTCCAACTCATCAATGT
-----
--SV40 sequences-->
3541  ATCTTATCATGTCTGGATCGGGGATCC

```

- 32 -

Insertion of the VV 7.5K promoter/lacZ gene construct
into FPV NERs to make recombination plasmids

05 The P7.5+lacZ fragment (about 3.6 kb) in 20 μ l total volume of
buffer was cut out of the ampicillin - resistant plasmid pPG1
with 10 units of BamHI. The ends of the DNA were then repaired
with the Klenow fragment of DNA polymerase to produce blunt-ended
DNA molecules. Thus, the fragment was phenol-extracted with an
equal volume of TE- saturated phenol and ethanol- precipitated.
It was resuspended in 20 μ l of ligation buffer containing 5 units
10 of Klenow fragment in 1 unit of T₄ polymerase, containing 0.025
mM of each of dATP, dCTP, dTTP and dGTP and stood for 1 hour at
room temperature. Similarly, the pB3ME (SpeI, NcoI and BglII)
were also digested with BamII and repaired to produce blunt-ended
sites. The P7.5+lacZ fragment was then blunt-end ligated into
15 pB3ME as follows. Thus, the plasmid and fragment (20 ng each)
were ligated in a total volume of 10 μ l above described ligation
buffer containing 1 unit of T₄ DNA ligase and 1mM ATP and
incubated at 4°C overnight and were transformed into E. coli.
Blue colonies of X-gal Kanamycin plates were selected (pBGS 19 is
20 a Kanamycin - resistant vector) and the DNA of several colonies
was examined by the method of Holmes and Quigley, loc. cit. Each
ligation produced two different results, i.e. the P7.5+lacZ could
be ligated into pB3ME in either of two different orientations.
The pairs of constructs of each site were called pEFL2/4, pEFL5/6
25 and pEFL7/8. In pEFL, 2, 6 and 8, the P7.5 + lacZ insert was in
the same orientation as the gene into which it was inserted i.e.
left to right in pEFL2 (BglII, 26.0) and pEFL8 (SpeI, 11.5) and
right to left in pEFL6 (NcoI, 13.9).

Recombination

30 The plasmids pEFL 2, 6 and 8 which contained the P7.5K
promoter/lacZ construct in the orientations described were used
in recombination experiments. The recombination has the
following general stages:

- 33 -

1. Infect CEF cells with FPV, and add the recombination plasmid DNA in the form of a pre-prepared calcium phosphate precipitate. Allow to grow for several days. (Recombination takes place)
- 05 2. Harvest cells and freeze-thaw three times (Virus is released from cells).
3. Add this preparation to CEFs at various dilutions to produce plates with about 500-1000 plaques per 10cm dish (about 0.1% of these are recombinant plaques).
- 10 4. Add Xgal overlay and visualise blue plaques.
5. Pick isolated plaques. Freeze-thaw three times to release virus from cells of plaque.
6. Titrate again on CEFs and get more plaques.
7. Loop back to 4, repeating stages 4-6, until a pure plaque of
- 15 the recombinant is obtained.

In more detail, chick embryo fibroblast cells, when about 80% confluent, were infected with about 3 plaque forming units (infectious virus particles) per cell of plaque-purified HP 440. fowlpoxvirus. The virus was contained in 1ml of commercially

20 available serum free medium ("199"), and was left to adsorb to the cells for 2 hours at 37°C, after which the virus inoculum was removed and 10ml of fresh medium containing 5% newborn calf serum (CS) was added. 1.5 hours later a precipitate of the plasmid DNA construct to be recombined into the virus was prepared: to 1ml of

25 HEPES-buffered saline (pH7.12) was added 10-20 µg of the pEFL 2, 6 or 8 plasmid DNA in solution in TE at about 1µg/µl. 50µl of 2.5M calcium chloride was added slowly, and a fine DNA/calcium phosphate precipitate was allowed to form for 30 minutes at room

30 temperature (25°C). The medium from the virus-infected cells was then removed, the 1ml of precipitate was added to the cells and left for 30 minutes at room temperature. Then 10ml of fresh medium containing 5% CS was added to the cells, and they were incubated at 37°C for a further 4 hours. The condition of the cells was then examined and if the DNA calcium phosphate

35 precipitate was considered too thick, the medium was replaced.

- 34 -

Otherwise it was then replaced the next day. The cells were incubated at 37°C for a further 5 days and then the virus was harvested by scraping the cells into the medium. Intracellular virus was released by three cycles of freeze-thawing and the virus titrated by a conventional plaque titration.

The liquid medium was removed from the cells and an overlay of cell culture medium in 1% low gelling temperature agarose (LGT) containing 300µg/ml of Xgal (5-bromo-4-chloro-3-indolyl-beta D-galactoside) was added. After overnight incubation at 37°C, plaques expressing the lacZ gene, blue in colour, were observed.

Blue plaques were picked into 500 µl of 10mM Tris pH9, intracellular virus was released by three cycles of freeze-thawing and the virus titrated by a conventional plaque titration. The process of picking plaques and re-titrating was repeated until a plaque purified recombinant was obtained.

The plaque-purified recombinants were then passaged in tissue culture for the following numbers of passages after the first picking:

20

recombinants from plasmid pEFL 8 (SpeI insertion): stable for at least 3 passages

6 (NcoI insertion): stable for only 2 passages

2 (BglII insertion): stable for at least 6 passages.

It is possible that the NcoI-based recombinants might be stable for a longer period under different conditions from those tried above.

- 35 -

DNA analysis of FPV recombinants

When the DNA of the FPV parent is digested with BamHI a number of restriction fragments are generated, which can be separated on agarose gels. The terminal fragment is of size 6.3kb and is of double intensity compared with the other bands since there are two copies of it (one from each end of the genome). When the DNA of the recombinant made with pEFL2 was digested the pattern of bands was the same except that the double 6.3kb fragment had been replaced by a double 9.9kb fragment. This is consistent with the increase in size from 6.3kb to 9.9kb which would be produced if the 3.6kb P7.5+lacZ fragment had been inserted into the terminal fragment. Probing of the restriction digest after transfer to nitrocellulose showed that the 9.9kb band did in fact contain lacZ sequences.

15

EXAMPLE 2

This Example relates to D8-D9 NERs.

In order to test whether D9, D10 and the fowlpox gene present in the D8 position were non-essential for growth further manipulations were carried out.

1. For D8 the plasmid pMB389 was grown in a dam⁻ E. coli host WK262 and reisolated as pMB434. The reason for this is that in the usual strains of E. coli, the enzyme BclI to be used here cannot restrict because its control adenine base in its TGATCA recognition sequence is methylated. The dam⁻ strain resists methylation and in this host the enzyme BclI would be able to cleave its two recognition sites which occurred in the D8 region. This plasmid was then cleaved with BclI and religated and a clone from which the 583 bp BclI fragment was missing was isolated by screening a number of colonies with small plasmid preps. Such a plasmid pMB441 containing a unique BclI site and having much of its D8 region removed was isolated. A BamHI fragment from pBPG1, containing the β -galactosidase gene, under the control of the vaccinia P7.5K promoter, was then inserted into this unique BclI site of pMB441 (BamHI and BclI generate compatible sticky ends such that no end-repairing was

- 36 -

necessary). A plasmid clone pMB447 containing the correct structure was isolated and used in recombination assays (see later).

05 2. For D9 pMB389 was cleaved with AsuII (a unique site for which exists in the D9 gene) and end-repaired. pBPGI was cleaved with BamHI and end-repaired and then ligated with AsuII cleaved pMB389 which had also been end-repaired. One colony selected as blue on Xgal Amp plates was found to contain the vaccinia P7.5K promoter and β -galactosidase gene inserted in the D9 gene.

10 3. For D10 an EcoRI fragment was subcloned from pMB389 (one EcoRI site lies within the D9 gene whilst the other is in the polylinker of the pUC13 vector of pMB389) into pUC13 cut with EcoRI. The resulting plasmid pMB439 contains a unique BglII site within the D10 gene. The BamHI fragment containing the vaccinia
15 P7.5K promoter and the β -galactosidase gene were cloned into the BglII site of pMB439 to generate pMB443. (Again no end-repair was necessary as BamHI and BglII also have compatible sticky ends). Plasmid pMB443 was then tested in recombination assays.

The plasmid constructions described above are illustrated
20 schematically in Fig. 10. The plasmid pBG1 is derived by cutting the BamHI fragment from pPG1, this fragment containing the P7.5- β -galactosidase gene construct, and inserting it in the plasmid pBGS19 (refer to Example 1).

Recombination

25 Viruses from initial blue plaques from a recombination experiment were plaque-purified three times and appeared stable (i.e. no white plaques were observed on the third passage). Viral DNA was made from the plaque-purified viruses, and digested with EcoRI. DNA from the original (non-recombinant) ("wild
30 type") FPV was also digested with EcoRI to serve as a control. The DNAs were run on duplicate agarose gels, Southern blotted, and probed with either the putatively recombinant FPV EcoRI fragment, within which the insertions should have taken place, or with pPG1 containing the β -galactosidase gene. Comparing the
35 putatively recombinant FPV with the wild type when the EcoRI

- 37 -

digests were run on gels, the normal EcoRI fragment present in wild-type DNA had increased in size in both the D8 and D9 putative recombinants. The new D8 fragment was smaller than the new D9 one, as expected, due to the removal of the small BclI fragment (377-960) in this case. When probed with β -galactosidase gene-containing pBG1 plasmid DNA, wild-type FPV DNA did not light up but the D8 and D9 putative recombinants both lit up in the same positions as when the probe from the putatively recombinant EcoRI fragment was used. This established that the structures of the D8 and D9 recombinants were as expected.

Similar experiments for the D10 gene failed to give the desired confirmations.

EXAMPLE 3

15 Virus strain.

This Example describes the construction of vectors containing Newcastle Disease Virus (NDV) F and HN genes, successful protection of birds using the F. gene. TIR NERs were used.

20 A. Subcloning of NDV Fusion (F) and Haemagglutinin-Neuraminidase (HN) genes.

The F and HN genes of NDV were provided as a gift from Professor P.T. Emmerson of the University of Newcastle-upon-Tyne, each cloned into the plasmid pUC19, P. Chambers et al., Journal of General Virology, 67, 2685-2694 (1986); N.S. Millar et al., Journal of General Virology, 67, 1917-1927 (1986). See also European Patent Application 227414 (National Research Development Corporation) which refers to Budapest Treaty patent deposits of NDV cDNA in plasmids pUC18, and pUC19 at the National Collection of Industrial Bacteria, Aberdeen, Scotland on 1st July 1986 as NCIB 12277 (F gene) and 12278 (HN gene). Referring to the pUC19 clones actually used herein, the F gene clone also contained part of the HN gene downstream thereof. The F gene was subcloned by cutting with the restriction enzyme BamHI (which conveniently cuts 29 bases upstream of the initiating ATG codon at the 5' end

- 38 -

of the F gene) and HindIII (which cuts approximately 820 bases downstream from the 3' end of the F gene, this extra sequence including some of the HN gene) into the kanamycin-resistant plasmid pBGS18 cut with BamHI and HindIII. The resulting plasmid was called pBGF1 (see Figure 11).

The pUC19 plasmid containing the HN gene also has F gene sequence upstream of the HN gene. In order to facilitate positioning of a poxvirus promoter adjacent to the HN gene, the F gene sequences were removed by the use of the exonuclease Bal31. The plasmid was cleaved with the restriction enzyme SphI (which cuts in the multiple cloning site of pUC19 on the 5' side of the HN gene) and digested for a range of appropriate times as follows. The method used was as described in "Molecular Cloning (A Laboratory Manual)" by T. Maniatis *et al.*, Cold Spring Harbor USA (1982). 20µl of DNA (at approximately 500µg/ml) were mixed with 20µl BSA (1mg/ml) and 40µl of 2 x "Bal31 buffer" (24mM CaCl₂, 24mM MgCl₂, 0.4M NaCl, 40mM Tris/HCl pH 8.0, 2mM EDTA) and warmed to 30°C. 0.5 unit of Bal31 was added and samples were incubated for 0, 1, 2, 3, 4 and 5 minutes. Then 20µl of water and 3µl of 0.5M Tris/HCl pH 7.5, 100mM MgCl₂ were added, and the samples were digested with 10 units of SstI (which cuts downstream of the HN gene) for 60 minutes. Samples were run on an agarose gel to estimate the amount of DNA digested, and the 0, 1 and 2 minute samples were chosen for further use. The molecules were then repaired (as described above) with T4 DNA polymerase and Klenow fragment and cloned into SmaI-cut pBGS18. Several different isolates were sequenced at the 5' end of the HN gene and one was found in which only 49 of the bases upstream of the gene remained (starting at the sequence GGCTTCACAA...). This plasmid was called pBHN1 (see Figure 13).

B. Cloning of the F gene into a recombination vector.

In order to place the F gene under the control of the P7.5K VV promoter, the F gene was excised from pBGF1 with BamHI and HindIII, end-repaired as described, and cloned into the SmaI site

- 39 -

of pGS20 (Mackett et al., Journal of Virology 49, 857-864). This plasmid was called pGSF1 (see Figure 11). A plasmid pEFL16 (Figure 12) was constructed by inserting the PvuII fragment from pUC19 into the SpeI site of pB3ME (described above). This
05 fragment contains the coding sequences of the alpha peptide of beta-galactosidase with a multiple cloning site arranged such that when fragments are cloned into this site the translation of the beta-galactosidase is disrupted and thus recombinants can be detected because they are white colonies on plates containing
10 Xgal. (This construct pEFL16 is now equivalent to the 'recombination plasmid' described in Example 1 and shown in Figure 7). The F gene plus P7.5 promoter was excised from pGSF1 by using XbaI and NheI, end-repaired as described, and cloned into the unique SmaI site of pEFL16. The resulting plasmid (which
15 now has the F gene, with the P7.5 promoter upstream, placed within the SpeI non-essential region of the terminal fragment of fowlpox) is called pEFF2 (Figure 12).

C. Cloning of the HN gene into a recombination vector.

20 In order to place the HN gene under the control of the P7.5 promoter it was excised from pBHN1 with SstI and BamHI, repaired as described, and cloned into the SmaI site of pGS20. This plasmid was called pGSHN1 (Figure 13). The HN gene plus P7.5 promoter was excised from pGSHN1 by using SalI and Asp718,
25 repaired as described, and cloned into pB3ME which had been cut with SpeI and repaired. This plasmid was called pEFH3 (see Figure 14).

D. Production of FPV recombinant containing the F gene.

30 The Poxine strain of fowlpox virus used was kindly donated by Duphar. It was passaged three times in chicken embryo fibroblast cells CEFs in tissue culture. A single plaque was picked at this stage and was plaque-purified three times. The virus thus obtained was called Px4.1. and was grown in tissue culture as
35 described for the HP444 strain in Example 1.

- 40 -

The plasmid pEFF2 was allowed to recombine into the Px4.1 strain of FPV, as described in Example 1. The virus from the recombinations was titrated on CEFs and then plaqued out on 10cm petri dishes at a concentration of virus producing about 1000
05 plaques per dish. The virus was overlaid with cell culture medium in 1% low gelling temperature agarose. After 4-5 days the agarose overlay was removed and stored at 4°C in a sterile petri dish. A dry nitrocellulose filter was laid onto the cell sheet and pressed down with a circle of Whatman 3MM chromatography paper soaked in 20 X SSC. The filter was then lifted off the
10 dish, with the cell sheet and plaques adhered to it, the 3MMs paper removed, and the filter was baked at 80°C in a vacuum for 2 hours. To identify recombinant viruses, the filters were then probed with radiolabelled probes specific for either the F gene or the HN gene. The filters were exposed to X-ray film and the
15 position of recombinant plaques identified by aligning the autoradiograms with the stored agarose overlays. Recombinant virus was then isolated from the agarose overlay by removing a plug of agarose using a pasteur pipette. This plug was resuspended in 10mM Tris pH 9.0, freeze-thawed three times and
20 replaques. Recombinant virus was identified again by probing, and this process was repeated three times in all until a plaque-purified virus was achieved.

E. Production of FPV recombinant containing the HN gene.

25 The plasmid pEFH3 was allowed to recombine, as described above, into the plaque-purified HP440 virus (HP 438 as described in Example 1, plus two further passages). This virus is called FP9. Recombinant virus containing the HN gene was isolated as described for the F gene.

30

F. Expression of the F gene in tissue culture.

The recombinant viruses, and the equivalent parent strain as controls, were used to infect 5cm petri dishes of CEFs at approximately 5 pfu/cell. The virus was allowed to grow for
35 various times and then the cells lysed by addition of 500µl of

- 41 -

boiling sample buffer (containing 24mM Tris/HCl pH 6.8, 1% SDS, 20% glycerol and 0.02% bromophenol blue). For the HN recombinant this buffer also included 0.1M dithiothreitol. The infected cells were scraped into the sample buffer, and the samples boiled
05 for 2 minutes, cooled on ice, and frozen at -20°C. Samples were then loaded onto standard polyacrylamide protein gels, electrophoresed to separate the proteins, and transferred to nitrocellulose filters by Western blotting. The Western blots were probed (as described in A.C.R. Samson et al., Journal of General Virology 67, 1199-1203) with a monoclonal antibody
10 specific to the F protein or the HN protein did not reveal any F protein produced by the recombinant. The F protein produced by infection of CEFs with NDV was detected by the monoclonal antibody, but at a lower level than the HN, and so if the same ratio of native to recombinant protein held for the F as for the
15 HN, the recombinant F might indeed not be seen.

G. Protection of birds against NDV challenge.

14 day old Rhode Island Red chickens were inoculated intravenously with 10^6 pfu/bird of the F gene recombinant (4
20 birds) or of a fowlpox strain FP9 having no inserted genes (6 birds). 6 birds were left uninoculated. The 10 birds not inoculated with the recombinant are hereinafter referred to as 'the control birds'. At 28 days the inoculation was repeated. At 64 days the birds were challenged intramuscularly with 10^5 x
25 ELD₅₀ (ELD = embryo lethal dose) of the virulent Herts 33 strain of NDV. At 3 days after challenge, 4 of the control birds were dead and 8 were very sick. At 4 days after challenge, 10 out of 12 control birds were dead, and by 5 days after challenge all the control birds were dead. All 4 birds inoculated with the F
30 recombinant were normal.

Serum was taken from birds inoculated with the F recombinant and control birds before challenge with NDV. Examination of this serum by probing Western blots of purified NDV virions, showed that only the serum from the birds inoculated with the F
35 recombinant had antibodies to the F protein. Thus although it

- 42 -

was not possible to detect F protein produced by the recombinant in vitro, the in vivo experiment showed very convincingly that protection could be achieved, and that F protein in some form has been produced by the FPV recombinant containing the NDV F gene.

05 Example 6 describes a protection experiment using the NDV HN gene constructs of this Example.

EXAMPLE 4

This Example demonstrates vaccination of birds at one day of age with an NDV F gene/ FPV recombinant.

10 Groups of six or seven Rhode Island Red chicks were inoculated, at one day of age, by the wing-web method with either the NDV F gene/poxine FPV recombinant described in Example 3 8907374 or the Px4.1 parent poxine FPV strain. Each virus strain was inoculated into four groups of birds, at doses of 5×10^4 ,
15 5×10^3 , 5×10^2 , and 5×10^1 pfu/bird. Thus there were eight groups in all, with an additional group of uninoculated birds. Birds were inoculated at one day of age and challenged intramuscularly 26 days later, as in Application 8907374. Table 1 shows the results of this challenge. Uninoculated and control
20 birds were not protected against challenge, all 38 birds being dead by 4 days. The birds inoculated with the fowlpox/NDV F recombinant were protected to a significant degree, 22 birds out of 28 surviving the challenge (79%). The dose of vaccine used appeared to have little effect on the levels of protection,
25 although the lowest dose gave the best protection.

- 43 -

TABLE 1

<u>Inoculum</u>	Viral <u>dose</u>	Total number of birds dead at days 3-6 after challenge with NDV Hert's 33 strain			
		days 3	day 4	day 5	day 6
NDV F gene/ Px4.1 FPV recombinant	5×10^4	0/7	0/7	2/7	2/7
	5×10^3	0/7	0/7	2/7	2/7
	5×10^2	1/7	1/7	2/7	2/7
	5×10	0/7	0/7	0/7	0/7
Px4.1 FPV alone	5×10^4	6/7	7/7		
	5×10^3	3/6	6/6		
	5×10^2	5/6	6/6		
	5×10	1/5	5/5		
Uninoculated	-	4/7	7/7		

EXAMPLE 5

This Example confirms the results given in Example 3 (Section G) for vaccination of birds with an NDV F gene recombinant and shows that it can be done by using the wing web as well as the intravenous route.

The procedure of Example 3 was followed, except that the booster inoculation was at 27 days from birth (rather than 28) and the challenge at 42 days from birth (rather than 64), the parent poxine strain was used as control, and there were five groups of birds (two inoculated iv, two wing web and one uninoculated).

The results in Table 2 show that all the birds inoculated with the NDV F gene/poxine FPV recombinant survived, whereas all the others died.

- 44 -

TABLE 2

<u>Inoculum</u>	<u>Method</u>	Total number of birds dead at days 3, 4 and 14 after challenge with NDV Herts 33 strain		
		day 3	day 4	day 14
NDV F gene/ Px4.1 FPV				
recombinant	iv	0/6	0/6	0/6
Px4.1 FPV alone	iv	6/6		
NDV F gene/ Px4.1 FPV				
recombinant	wing-web	0/6	0/6	0/6
Px4.1 FPV alone	wing-web	4/6	6/6	
Uninoculated	-	9/13	13/13	

EXAMPLE 6

This Example shows that a NDV HN gene/FPV recombinant also gives protection. Thus, protection can be obtained from either the F or the HN gene alone.

05 Example 5 was repeated except that the NDV HN/HP 440 FPV recombinant as described in Example 3 (Section E) was used with the HP 440 strain as control.

Table 3 shows the results. Again, the NDV HN gene-vaccinated birds were protected while the others all died.

TABLE 3

<u>Inoculum</u>	<u>Method</u>	Total number of birds dead at days 3, 4 and 14 after challenge with NDV Herts 33 strain		
		day 3	day 4	day 14
NDV HN gene/ HP 440 FPV				
recombinant	iv	1/6	1/6	1/6
HP 440 alone	iv	6/6		
Uninoculated	-	9/13	13/13	

- 45 -

EXAMPLE 7

In this Example recombination vectors were made using the Ecogpt gene, the constructs being of a generally similar kind to those described by CSIRO in the already cited PCT Application
05 WO 88/02022, but flanked by non-essential region sequences of the present invention rather than TK gene sequences. The recombination vectors comprise in order:

- (1) Non-essential region (around the BglII site)
- (2) Ecogpt gene
- 10 (3) VV P.5 promoter
- (4) VV P11 promoter
- (5) IBV spike protein gene
- (6) Non-essential region.

Note that the two promoters are "back-to-back", as shown in
15 WO 88/02022, transcribing in opposite direction.

The clone pEFL2 (containing the vaccinia P7.5 promoter and beta-galactosidase gene in the FPV terminal fragment, see above) was cut with the restriction enzymes SphI and NcoI, repaired as described, and religated. The resulting construct, designated
20 pEFL18, had lost approximately 850 base pairs of FPV sequence, as expected, but had also lost a HindIII site immediately adjacent to the SphI site. Thus, this plasmid contains only two HindIII sites, very close together, between the P7.5 promoter and the beta-galactosidase gene. (Digestion with HindIII thus only
25 removes a very small fragment and can be used for the insertion of gene fragments containing their own ATG codon to make beta-galactosidase fusion constructs). This plasmid was cut with HindIII and EspI (EspI cuts near to the 3' end of the beta-galactosidase gene) and repaired. Into this repaired site
30 was inserted a BglII/ApaI fragment (repaired) from the plasmid pSV2gpt (Mulligan & Berg, Science 209, 422-427, 1980) which contained the E.coli ecogpt gene. Thus, the beta-galactosidase gene was removed and the Ecogpt gene was placed under the control of the P7.5 promoter. A unique SalI site upstream of the
35 promoter allows insertion of other promoters or promoter/gene constructs. This plasmid was called pEFGP13.

- 46 -

The P11 late vaccinia promoter was excised from pMM6 (a gift from Michael Mackett) provided with multiple cloning sites (BamHI, XhoI, XbaI and SalI) by transfer to pUC19 and the resultant P11-cloning site construct inserted into the SalI site of pEFGPT3. The resultant plasmid was designated pEFGPT6 and allows insertion of genes at various sites under the control of the P11 promoter. An Infectious Bronchitis Virus spike protein gene (see NRDC's PCT Application Publication No. WO86/05806) was inserted downstream of the promoter.

10 The plasmid pEFGPT3 was allowed to recombine into FPV strain FP9, as described above. Recombinant viruses were selected by the use of MXHAT medium containing mycophenolic acid (Boyle & Coupar, Gene 65, 123-129, 1988). After the initial recombination in the presence of MXHAT medium, the virus was released by
15 freeze-thawing and passaged twice under selective conditions. A cytopathic effect was seen, indicating that virus was growing under the selective conditions. Virus was then plaqued out and probed with an EcoGpt-specific probe. A high proportion of plaques (approximately 75%) were positive, indicating that the
20 selection had been successful (normally only 0.1% of the virus would be recombinants) and hence that recombinants expressing the EcoGpt gene had been isolated.

Similar recombination experiments were carried out to incorporate the EcoGpt gene into the Poxine strain of FPV. After
25 the recombination and upon passage in selective conditions, a cytopathic effect of the virus could be seen, indicating that recombinant virus expressing the EcoGpt gene had been produced. This confirms that the BglII non-essential region (as well as the SpaI region) is present in Poxine.

- 47 -

CLAIMS

1. A recombination vector for use in producing a recombinant fowlpox virus (FPV) by homologous DNA recombination, said recombination vector comprising a cloning vector containing a non-essential region (NER) sequence comprising NER of (a) at
05 least part of the long unique sequence (LUS) of the terminal inverted repeat (TIR) of FPV or (b) at least part of the region of the FPV genome lying between the carboxy-end of the D7 gene and the carboxy-end of the D9 gene (but excepting the D10 gene and D10 gene promoter region), said NER being interrupted by at
10 least one foreign gene and by at least one sequence for regulating the expression of the or each foreign gene.
2. A recombination vector according to Claim 1 wherein the NER is wholly within the LUS of the TIR.
3. A recombination vector according to Claim 2 wherein the NER
15 lies wholly within 2.2 kilobase pairs from the external end of the LUS.
4. A recombination vector according to Claim 3 wherein the NER lies within an open-reading frame (ORF) which is substantially one of those at 4630-4695, 5166-5534 or 5788-6090 and having the
20 nucleic acid sequence herein identified or a variant thereof occurring in another strain of FPV.
5. A recombination vector according to Claim 1 wherein the NER lies within the D8 or D9 gene.
6. A recombination vector according to Claim 1, which comprises
25 in order :
 - (1) a first homologously recombinable flanking sequence,
 - (2) a sequence within a first portion of the non-essential region,
 - (3) promoter DNA,
 - 30 (4) a foreign gene transcribably downstream of the promoter (whereby when the fowlpox virus RNA polymerase binds to the promoter it will transcribe the foreign gene into mRNA),

- 48 -

- (5) a sequence within a second portion of the same non-essential region, the first and second sequences being in the same relative orientation as are the first and second portions of the non-essential region within the viral genome, and
- 05 (6) a second homologously recombinable flanking sequence.
7. A recombinant vector useful in preparing a recombination vector according to Claim 1, this recombinant vector containing a NER sequence as defined in Claim 1,2,3,4 or 5.
- 10 8. A DNA molecule which comprises a NER sequence as defined in Claim 1,2,3,4 or 5, with or without up to 1000 bp of homologously recombinable flanking sequence at either or both ends thereof.
9. A recombinant fowlpox virus (FPV) which is the product of homologous recombination of a parent FPV with the insert DNA of a recombination vector claimed in any one of Claims 1-6.
- 15 10. An in vitro culture of animal cells infected with a recombinant FPV according to Claim 9.
11. A method of vaccinating an animal, which comprises inoculating it with a recombinant FPV as defined in Claim 9 or a culture according to Claim 10.
- 20 12. A method according to Claim 11 wherein the animal is a chicken.

1 / 14





Fig. 1



2 / 14





-  *Short unique sequence*
-  *Tandemly repeated sequences*
-  *Beginning of the long unique sequence*
-  *Vector sequences*

Fig. 2















Fig. 3



-  *Foreign gene*
-  *Promoter*
-  *Non-essential region*
-  *Plasmid DNA*



-  *Foreign gene*
-  *Promoter*
-  *Non-essential region*
-  *Multiple cloning site*
-  *Plasmid DNA*




Fig. 6


Foreign gene for avian antigen

Foreign marker gene

Promoters

Non-essential region

Plasmid DNA



Foreign gene for avian antigen


Foreign marker gene

Promoters

Non-essential region

Multiple cloning site

Plasmid DNA






Fig. 9




Fig. 10

11 / 14




Fig. 11

12 / 14




Fig.12

13 / 14




Fig. 13

14 / 14




Fig. 14

INTERNATIONAL SEARCH REPORT

International Application No PCT/GB 89/00698

I. CLASSIFICATION OF SUBJECT MATTER (if several classification symbols apply, indicate all) ⁶		
According to International Patent Classification (IPC) or to both National Classification and IPC		
IPC ⁴ : C 12 N 15/00, C 12 N 7/00, C 12 N 5/00		
II. FIELDS SEARCHED		
Minimum Documentation Searched ⁷		
Classification System	Classification Symbols	
IPC ⁴	C 12 N	
Documentation Searched other than Minimum Documentation to the extent that such Documents are included in the Fields Searched ⁸		
III. DOCUMENTS CONSIDERED TO BE RELEVANT ⁹		
Category ⁹	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³
X,P	EP, A, 0284416 (NIPPON ZEON CO. LTD) 28 September 1988, see example 2 --	1-10
X,P	WO, A, 89/03429 (HEALTH RESEARCH INC.) 20 April 1989, see page 16 cited in the application --	1-10
X,P	EP, A, 0308220 (NIPPON ZEON CO. LTD) 22 March 1989, see page 4, lines 54-61; figure 6 --	1-10
A	WO, A, 86/00528 (GENEX CO.) 30 January 1986, see page 26, lines 6-24 -----	1-10
<div style="display: flex; justify-content: space-between;"> <div style="width: 45%;"> <p>¹⁰ Special categories of cited documents:</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 45%;"> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"A" document member of the same patent family</p> </div> </div>		
IV. CERTIFICATION		
Date of the Actual Completion of the International Search	Date of Mailing of this International Search Report	
29th September 1989	27 OCT 1989	
International Searching Authority	Signature of Authorized Officer	
EUROPEAN PATENT OFFICE	T.K. WILLIS	

FURTHER INFORMATION CONTINUED FROM THE SECOND SHEET

V. ☒ OBSERVATIONS WHERE CERTAIN CLAIMS WERE FOUND UNSEARCHABLE ¹

This international search report has not been established in respect of certain claims under Article 17(2) (a) for the following reasons:

1. ☒ Claim numbers* because they relate to subject matter not required to be searched by this Authority, namely:

* 11,12

See PCT Rule 39.1(iv): Methods for treatment of the human or animal body by surgery or therapy, as well as diagnostic methods

2. ☐ Claim numbers because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. ☐ Claim numbers because they are dependent claims and are not drafted in accordance with the second and third sentences of PCT Rule 6.4(a).

VI. ☐ OBSERVATIONS WHERE UNITY OF INVENTION IS LACKING ²

This International Searching Authority found multiple inventions in this international application as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims of the international application.
2. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims of the international application for which fees were paid, specifically claims:
3. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claim numbers:
4. ☐ As all searchable claims could be searched without effort justifying an additional fee, the International Searching Authority did not invite payment of any additional fee.

Remark on Protest

- ☐ The additional search fees were accompanied by applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

**ANNEX TO THE INTERNATIONAL SEARCH REPORT
ON INTERNATIONAL PATENT APPLICATION NO.**

GB 8900698

SA 29725

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report. The members are as contained in the European Patent Office EDP file on 20/10/89
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A- 0284416	28-09-88	AU-A- 1376688	29-09-88
WO-A- 8903429	20-04-89	AU-A- 2427588	02-05-89
		FR-A- 2621487	14-04-89
		NL-A- 8820679	03-07-89
EP-A- 0308220	22-03-89	JP-A- 1157381	20-06-89
WO-A- 8600528	30-01-86	EP-A- 0190254	13-08-86
		JP-T- 61502586	13-11-86